





Committed to trials of LDN as a treatment for Autoimmune Disease

PO Box 1083, Buxton, Norwich, NR10 5WY, UK.

Contact: Linda Elsegood  
Web: [www.ldnresearchtrust.org](http://www.ldnresearchtrust.org)  
email: [contact@ldnresearchtrust.org](mailto:contact@ldnresearchtrust.org)

Skype Phone No's

 01223 92 6933  
 (847) 794-8046

## Low-dose Naltrexone (LDN) Fact Sheet 2016

Naltrexone is in a class of drugs known as an opioid antagonists. Its normal use is in treating addiction to opiate drugs such as heroin or morphine. The dose used for this purpose is usually between 50 and 300mg daily.

Low-dose Naltrexone (LDN) has been used in the treatment of autoimmune diseases in the USA since 1985, but is relatively new in the United Kingdom and Europe. Despite the fact that the drug is used at a very low dose, the occurrence of significant introductory or long term side effects cannot be excluded.

This method was devised and subsequently developed by the late Dr. Bernard Bihari, a Neurophysician from New York, USA who passed away on May 16th 2010. Dr. Bihari was qualified in Internal Medicine, Psychiatry and Neurology, and we hope to honour him by continuing with his pioneering work.

### Suggested Method of Therapy

LDN has been used for over a decade internationally, so internet searches will bring up conflicting information about dosing and optimal time of day to take LDN. The LDN Research Trust produces this factsheet to inform clinicians and patients of the most up to date and practical uses of LDN.

In order to avoid common side effects, such as GI disturbance, headache, increased malaise, flu-like symptoms, agitation or worsening tiredness/spasticity in MS – doses should be started at 0.5-1mg and increased gradually.

The most common form of LDN is Liquid 1mg/1ml, and this can be started at 0.5mg for 7 days, increasing by 0.5mg weekly until at 3mg or 4.5mg, whichever is tolerated best.

As of July 2016, LDN 1mg/1ml from Dickson's Chemist in Glasgow is stable at room temperature for 12 months. Other suppliers do not follow this formula so liquid LDN from these sources may be stable for only 28 days. Many websites and internet resources state that LDN only works if taken at nighttime; however, clinical experience over the last decade clearly demonstrates that it is as effective in the morning – with far fewer side effects.

After starting LDN, some response is expected to be seen at 6 weeks, with maximal effects at 12 weeks.


## How Naltrexone Works

As of 2016, LDN is most commonly being used for Chronic Fatigue, Multiple Sclerosis, Myelagic Encephalopathy, autoimmune thyroid diseases and various cancers. Many autoimmune diseases seem to respond to LDN.

This is a wide range of diseases and many clinicians will find it difficult to understand how one drug can have a positive effect on all these pathologies.

The first thing to understand is that Naltrexone – the drug in LDN – comes in a 50:50 mixture of 2 different shapes (called isomers). It has been recently discovered that one particular shape binds to immune cells, whilst the other shape binds to opioid receptors.

Although consisting of exactly the same components, the two *isomers* appear to have different biological activity.



**The LEVO (left handed) version of naltrexone blocks opiate receptors.**

**The DEXTRO (right handed) version blocks receptors on immune cells. These include “Toll Like Receptors” which are heavily involved in immunity.**

For clinicians interested in reading more about the pharmacology behind this, there is a published resource available here:

**EU:** <https://www.amazon.co.uk/Ldn-Book-Little-Known-Naltrexone-Revolutionize/dp/1603586644>

**USA:** [https://www.amazon.com/LDN-Book-Little-Known-Naltrexone-Revolutionize/dp/1603586644/ref=sr\\_1\\_1?ie=UTF8&qid=1470137762&sr=8-1&keywords=ldn+book](https://www.amazon.com/LDN-Book-Little-Known-Naltrexone-Revolutionize/dp/1603586644/ref=sr_1_1?ie=UTF8&qid=1470137762&sr=8-1&keywords=ldn+book)

## Summary of mechanism of action

The summary of 10 years of research is that LDN works because:

Levo-Naltrexone is an antagonist for the opiate/endorphin receptors

- This causes increased endorphin release
- Increased endorphins modulate the immune response
- This reduces the speed of unwanted cells growing

Dextro-Naltrexone is an antagonist for at least one, if not more immune cells

- Antagonises "TLR," suppressing cytokine modulated immune system
- Antagonises TLR-mediated production of NF-kB – reducing inflammation, potentially downregulating oncogenes

Taking Naltrexone in larger doses of 50-300mg seems to negate the immunomodulatory effect by overwhelming the receptors, so for the effect to work, the dose must be in the range of 0.5-10mg, usually maxing at 4.5mg in clinical experience.

## The Use of Low-dose Naltrexone, and the Occurrence of Side Effects

Many patients who start LDN do not experience any severe side effects.

As mentioned earlier, your symptoms may become worse – in MS, this can be characterised by increased fatigue, or increased spasticity. In CFS/ME, this can be the onset of apparent flu-like symptoms.

LDN can cause sleep disturbances if taken at nighttime – this is most likely because of the increase in endorphin release. These disturbances can take the form of vivid dreams, or insomnia.

In various studies (and anecdotal accounts), the number of T-Lymphocytes has been shown to dramatically increase when a patient starts on LDN. This may account for some of the benefits patients feel when they are being treated for an autoimmune disease, or cancer. This has not been directly evidenced in multiple sclerosis.

Clinical experience shows that in less than ten percent of cases treated, increased introductory symptoms may be more severe or more prolonged than usual, lasting sometimes for several weeks. Rarely, symptoms may persist for two or three months before the appropriate beneficial response is achieved.

If side effects are troublesome, then reducing your dose by 50% for 7 days, before increasing it again, is a good idea.

Some patients, very rarely, experience gastro-intestinal side effects, such as nausea and or constipation/diarrhea. The reason for this is currently unknown, but may be due to the presence of large numbers of delta-opiate receptors in the intestines.

*Patients experiencing this side effect can request LDN Sublingual Drops, which transfer the LDN directly into the bloodstream – avoiding the GI tract.*

Patients who do have these side effects should increase their dose by no more than 0.5mg per week, and should consult with their GP or pharmacist for appropriate treatment for the stomach upset, if necessary. (Omeprazole, Ranitidine, Gaviscon, Fybogel, Mucogel and Pepto Bismol are ok – but not Kaolin & Morphine or Loperamide/Imodium.)

## Types of LDN

### Liquid

Oral Liquid Formulation at 1mg/1ml is the most commonly used type of LDN.

It is taken daily, and dosed using a baby oral syringe. It does not contain very high amounts of lactose or any other excipient known to cause hypersensitivity. The base is similar to children's cough syrup, and so is quite palatable. Because there are so few preservatives, it should be stored in the fridge. Cost: around 50p a dose and can last for 3 months or longer. The product no longer has to be refrigerated, but it is good practice to keep the product in the fridge.

### Capsules

For patients who find the liquid would be impractical, there are capsules available in 3mg and 4.5mg strengths in the UK. Other strengths are available from different manufacturers – but these are made on a case by case basis. These have up to 12 months stability data and can be stored anywhere. They contain no lactose filler and are instant release. Cost: Approx £1 a dose.

When asking a doctor to prescribe LDN, it is important to ask them to state “**LOWTREX - LDN**” as this will guarantee that you receive the same product, of high quality, and at the lowest available cost. This is available in 3mg and 4.5mg strengths.

### Sublingual Drops

Sublingual drops are designed for patients who are having problems taking the medication orally, or for people who want to guarantee the fastest delivery of the drug into their bloodstream. A number of drops are placed under the tongue from a dropper bottle and dose is increased and decreased by the number of drops taken. There are basically no excipients in this product, trace lactose and a small amount of glycerol. Cost about £1 a dose.

### Cream

LDN Cream in 0.5mg/ml is available for application to the skin. This is helpful for children, or for patients allergic to colourants, flavourants or any excipients in all other forms of LDN. It is more expensive at about £2 per dose, and lasts for 28 days only.

The efficacy of this product in adults is extremely dubious, as absorption into the bloodstream is very low.

*NB: Naltrexone has been investigated as a method of speeding up wound healing time – LDN Cream, unless specifically specified on the prescription, will not be sterile and cannot be used for this purpose. Check with your pharmacist first. Sterile cream will be significantly more expensive. Using non sterile cream on an open wound would be very dangerous.*

### **Intrinsic Toxicity of the Drug:**

Naltrexone, in full doses of 50-300mg, has been shown to transiently increase liver enzymes. Patients being prescribed Naltrexone for addictions must have liver function tests performed before initiating therapy.

This is not necessary with LDN as the dose is much smaller, however, patients with advanced liver failure should consult their GP before considering treatment.

Patients with renal or liver failure should only start treatment after a consultation with their own GP or specialist, and should be monitored during the treatment initiation period. It is normal for people with poor renal or liver function to experience a transient elevation – but

this usually resolves after a few weeks.

**Contraindications and Special Precautions:**

LDN is compatible with most other therapies. It does not directly interact with steroids, however, LDN can negate the effect of opiate based painkillers. Patients should give their doctor a full drug history before starting therapy.

Patients who are taking multiple medications and/or herbal medicines, especially those with cancer or advanced disease, should take careful advice from a qualified doctor or pharmacist before initiating LDN.

The LDN Research Trust retains medical and pharmaceutical advisors to assist with this process. If you are unsure whether you can take LDN with your current medications, please email us or our pharmaceutical advisor, Stephen Dickson MRPharmS. directly on:

[www.askaboutldn.com](http://www.askaboutldn.com)

<mailto:homedeliverypharmacy@yahoo.co.uk>

## **Low Dose Naltrexone - Key clinical studies**

Low Dose Naltrexone has been the subject of much debate but actually very few clinical trials. Ian Zagon from Penn State University has been studying LDN for over 20 years and conducted many pre-clinical studies investigating LDN in cancer and in the animal model of MS (1,2). He has also been involved in two clinical studies into Crohn's disease with his colleague Professor Jill Smith from Penn State. These demonstrated a significant improvement in symptoms and in bowel mucosal appearance with LDN treatment (3,4). In the RCT, LDN patients were twice as likely to have a 70 point decline in the Crohn's Disease Activity Index. 78% of the LDN group achieved an endoscopic response compared to 28% with placebo.

Jarred Younger from Stanford University has studied LDN in Fibromyalgia, firstly in a small pilot study and more recently in a yet to be published randomised controlled trial. The pilot study showed significant improvement in symptoms of pain in these patients (5).

Multiple Sclerosis is one of the areas where LDN has been used the most frequently. There are three published studies, one in primary progressive MS (6) and two on quality of life (7,8). The results of two studies was positive with improved quality of life in one and reduced spasm in the PPMS study. The third showed no significant difference between the treatment and placebo groups but found the treatment to be safe. A review of the available studies into LDN and MS was published in 2009 (9). All studies have confirmed the safety of the drug and there is enough positive evidence to merit greater investigation.

**Key references :**

1. Rahn KA, McLaughlin PJ, Zagon IS. Prevention and diminished expression of experimental autoimmune encephalomyelitis by low dose naltrexone (LDN) or opioid growth factor (OGF) for an extended period: Therapeutic implications for multiple sclerosis. Brain Res. 2011 Mar 24;1381:243-53. Epub 2011 Jan 20.
2. Donahue RN, McLaughlin PJ, Zagon IS. The opioid growth factor (OGF) and low dose naltrexone (LDN) suppress human ovarian cancer progression in mice. Gynecol Oncol. 2011 Aug;122(2):382-8. Epub 2011 Apr 30.

3. Smith JP, Stock H, Bingaman S, Mauger D, Rogosnitzky M, Zagon IS Low-dose naltrexone therapy improves active Crohn's disease. Am J Gastroenterol. 2007 Apr;102(4):820-8. Epub 2007 Jan 11.
  4. Smith JP, Bingaman SI, Ruggiero F, Mauger DT, Mukherjee A, McGovern CO, Zagon IS. Therapy with the opioid antagonist naltrexone promotes mucosal healing in active Crohn's disease: a randomized placebo-controlled trial. Dig Dis Sci. 2011 Jul;56(7):2088-97. Epub 2011 Mar 8.
  5. Younger J, Mackey S. Fibromyalgia symptoms are reduced by low-dose naltrexone: a pilot study. Pain Med. 2009 May-Jun;10(4):663-72. Epub 2009 Apr 22
  6. Gironi M, Martinelli-Boneschi F, Sacerdote P, Solaro C, Zaffaroni M, Cavarretta R, Moiola L, Bucello S, Radaelli M, Pilato V, Rodegher M, Cursi M, Franchi S, Martinelli V, Nemni R, Comi G, Martino G. A pilot trial of low-dose naltrexone in primary progressive multiple sclerosis. Mult Scler. 2008 Sep;14(8):1076-83
  7. Cree BA, Korniyeva E, Goodin DS Pilot trial of low-dose naltrexone and quality of life in multiple sclerosis. Ann Neurol. 2010 Aug;68(2):145-50.
  8. Sharafaddinzadeh N, Moghtaderi A, Kashipazha D, Majdinasab N, Shalbafan B. The effect of low-dose naltrexone on quality of life of patients with multiple sclerosis: a randomized placebo-controlled trial. Mult Scler. 2010 Aug;16(8):964-9. Epub 2010 Jun 9.
- Gilhooly TC Low-dose naltrexone as a treatment for multiple sclerosis  
British Journal of Neuroscience Nursing, Vol. 5, Iss. 11, 13 Nov 2009, pp 494

The latest up-to-date references are always being added to the website, however, the current authority is the LDN BOOK available on Amazon.

#### Obtaining a prescription for LDN:

**\*\*\*\*\*WARNING\*\*\* DO NOT buy LDN on the internet. There is no guarantee that the drug is genuine or safe. On multiple occasions LDN purchased from the internet or from overseas has been proven to be of low quality, completely fake or otherwise dangerous.**

The only way to legally and safely obtain LDN, is via a doctor's prescription.

#### NHS:

Take this factsheet first to your NHS GP. LDN will pass for payment on the NHS in Scotland, England, N.Ireland and Wales when written in the following manner:

**Liquid form**  
**Rx**

**NALTREXONE TABLETS 50mg**  
**SEND: 28 days supply**  
**DIRECTIONS:**

**EXTEMP PREP in pharmacy to 1mg/1ml. (ref Fawcett et al)**

**"Take Xxml daily for 7 days, then increase by 0.5mg a week until taking 4.5mg"**

## Capsule

Rx

**Lowtrex-LDN Capsules 3mg OR 4.5mg**

Send: 28 days' supply

Directions: One daily.

As LDN is unlicensed, your own GP may refuse to prescribe it (for insurance reasons), and you may need to go privately. **Lowtrex** is the most widely used brand in the UK and is made in the UK under "GMP" - their factory is inspected by the MHRA for safety and the product tested regularly for safety.

### PRIVATE:

**If your NHS GP is unable to prescribe you LDN, a list of currently prescribing private doctors is available at:**

[www.ldn-international.com](http://www.ldn-international.com) or via [www.askaboutldn.com](http://www.askaboutldn.com)

### Filling an LDN Prescription:

Prescriptions, whether NHS or Private, can be posted to:

Dickson Chemist Ltd  
LDN DEPT  
35 Mitchell Arcade  
Rutherglen  
Glasgow  
G73 2LS  
TEL: 0141 647 8032

to be dispensed.

email: [homedeliverypharmacy@yahooco.uk](mailto:homedeliverypharmacy@yahooco.uk)

website: [www.dicksonchemist.co.uk](http://www.dicksonchemist.co.uk) and [www.homedeliverypharmacy.co.uk](http://www.homedeliverypharmacy.co.uk)

The LDN is sent directly to your home – and is free if you have an NHS prescription.

If you are paying privately, liquid is around ~£18 for 28days supply, and capsules are ~£30 for 28 days supply.

Even if you have an NHS prescription, please consider sending it to Dickson's, as they only charge the NHS a minimal fee. Many other pharmacies may have it made specially and charge the NHS in excess of £150 for 28days supply.

You can also use any pharmacy of your choice – however, Dickson Chemist are by far the most experienced, least expensive and will send LDN to most EU and international

destinations.